


In the Claims:

1. (Currently Amended) A compound of the formula




wherein R, R₁ and R₂ are individually hydrogen or lower alkyl; X is

-O- or -NH-; PAG is a divalent residue of polyalkylene glycol resulting from removal of both of its terminal hydroxy groups, which residue has a molecular weight of from 1,000 to 50,000 Daltons; n is an integer of from 0 to 1; m is an integer of from 4 to 8; and A is a hydrogen or an activated leaving group which when taken together with its attached oxygen atom forms an ester

or hydrolyzable esters thereof wherein A is hydrogen, wherein said PAG residue has a molecular weight of about 10,000 to about 40,000 Daltons when X is O.

2. (Original) The compound of claim 1 having the formula




I-A1

wherein A, R, PAG, R¹, R², m and n are as above.

3. (Original) The compound of claim 2 wherein A is hydrogen.
4. (Original) The compound of claim 3 wherein PAG is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
5. (Original) The compound of claim 4 wherein R is methyl.
6. (Original) The compound of claim 5 wherein n is 0 and m is 4.
7. (Original) The compound of claim 5 wherein PEG has a molecular weight of from 10,000 to 40,000.
8. (Original) The compound of claim 6 wherein PEG has a molecular weight of from 20,000 to about 35,000.
9. (Original) The compound of claim 2 wherein A is an activated leaving group.
10. (Original) The compound of claim 9 wherein PAG is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
11. (Original) The compound of claim 9 wherein R is methyl.
12. (Original) The compound of claim 11 wherein n is 0 and m is 4.
13. (Original) The compound of claim 12 wherein PEG has a molecular weight of from 10,000 to 40,000.
14. (Original) The compound of claim 13 wherein PEG has a molecular weight of from 20,000 to about 35,000.

15. (Original) The compound of claim 1 wherein said compound has the formula




I-A2

wherein A, R, PAG, R¹, R², m and n are as above.

16. (Original) The compound of claim 15 wherein A is hydrogen.
17. (Original) The compound of claim 16 wherein PAG is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
18. (Original) The compound of claim 17 wherein R is methyl.
19. (Original) The compound of claim 18 wherein n is 0 and m is 4.
20. (Original) The compound of claim 19 wherein PEG has a molecular weight of from 10,000 to 40,000.
21. (Original) The compound of claim 20 wherein PEG has a molecular weight of from 20,000 to about 35,000.


22. (Original) The compound of claim 18 wherein PAG is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
23. (Original) The compound of claim 22 wherein R is methyl.
24. (Original) The compound of claim 23 wherein n is 0 and m is 4.
25. (Original) The compound of claim 24 wherein PEG has a molecular weight of from 10,000 to 40,000.
26. (Original) The compound of claim 25 wherein PEG has a molecular weight of from 20,000 to about 35,000.
27. (Original) The compound of formula



wherein R is hydrogen or lower alkyl; X is -O- or -NH-; PAG is a divalent residue of polyalkylene glycol resulting from removal of both of its terminal hydroxy groups, which residue has a molecular weight of from 1,000 to 50,000 Daltons ; w is an integer of from 1 to 3; and one of R₃ and R₄ is lower alkyl and the other is hydrogen or lower alkyl; and A is a hydrogen or an activated leaving group which when taken together with its attached oxygen forms an ester;

or hydrolyzable esters thereof wherein A is hydrogen.

28. (Original) The compound of claim 27 wherein said compound is




I-B1

wherein A, R, PAG, R^3 , R^4 , w and n are as above.

29. (Original) The compound of claim 28 wherein A is hydrogen.
30. (Original) The compound of claim 29 wherein PAG is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
31. (Original) The compound of claim 30 wherein R is methyl.
32. (Original) The compound of claim 31 wherein w is 1.
33. (Original) The compound of claim 32 wherein PEG has a molecular weight of from 10,000 to 40,000.
34. (Original) The compound of claim 33 wherein PEG has a molecular weight of from 20,000 to about 35,000.
35. (Original) The compound of claim 28 wherein A is an activated leaving group.
36. (Original) The compound of claim 35 wherein PAG is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
37. (Original) The compound of claim 36 wherein R is methyl.
38. (Original) The compound of claim 37 wherein w is 1.
39. (Original) The compound of claim 38 wherein PEG has a molecular weight of from 10,000 to 40,000.

40. (Original) The compound of claim 39 wherein PEG has a molecular weight of from 20,000 to about 35,000.


41. (Original) The compound of formula



wherein R is hydrogen or lower alkyl, X is -O- or -NH-, A is a hydrogen or an activated leaving group which when taken together with its attached oxygen atom forms an ester, PAG¹ is a divalent residue of a polyalkylene glycol resulting from the removal of both of the terminal hydroxy groups, said residue having a molecular weight of from about 500 to about 25,000 Daltons, y is an integer from 0 to 3 and v is an integer from 1 to 3; and k is an integer from 1 to 2;

or hydrolyzable esters thereof wherein A is hydrogen.

42. (Original) The compound of claim 41 wherein said compound has the formula




wherein R, PAG¹, A v, y and k are all as above.

43. (Original) The compound of claim 42 wherein A is hydrogen.

44. (Original) The compound of claim 43 wherein PAG¹ is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.

45. (Original) The compound of claim 42 wherein each PAG¹ residue has a molecular weight of 500 to 15,000.
46. (Original) The compound of claim 42 wherein A is a leaving group.
47. (Original) The compound of claim 46 wherein said leaving group is N-hydroxysuccinimidyl.
48. (Original) The compound of claim 47 wherein PAG¹ is PEG, a divalent polyethylene glycol residue resulting from the removal of both of its terminal hydroxy groups.
49. (Original) The compound of claim 48 wherein R is methyl.
50. (Original) The compound of claim 49 wherein each PEG residue has a molecular weight of from 500 to 10,000.
51. (Currently Amended) A process for producing an activated ester of the formula:



wherein R, R₁ and R₂ are individually hydrogen or lower alkyl; X is -O- or -NH-; PAG is a divalent residue of polyalkylene glycol resulting from removal of both of its terminal hydroxy groups, which residue has a molecular weight of from 1,000 to 50,000 Daltons; n is an integer of from 0 to 1; m is an integer of from 4 to 8; and A is a hydrogen or an activated

leaving group which when taken together with its attached oxygen atom
 forms an ester


comprising, condensing a compound of the formula:



V

wherein R, and PAG are as above, and V is $-\text{OH}$ or $-\text{NH}_2$,


with the compound of the formula:



wherein R^5 forms a hydrolyzable ester protecting group and Y is halide and


R^1 , R^2 , m, and n, are as above,

to produce an ester of the formula



wherein R, PAG, X, R^1 , R^2 , R^5 , m and n are as above,

hydrolyzing said ester to form a free acid of the formula:



wherein R, PAG, X, R¹, R², m and n are as above,


and reacting said free acid with a halide of an activated leaving group in the presence of a coupling agent to produce said activated ester,

and wherein said PAG residue has a molecular weight of about 10,000 to about 40,000


Daltons when X is O.

52. (Currently Amended) The process of claim 51 wherein said leaving group is a N-hydroxysuccinimidyl group~~58.~~

53. (Original) A process for producing an activated ester of the formula:



wherein R is hydrogen or lower alkyl; X is -O- or -NH-; PAG is a divalent residue of polyalkyleneglycol resulting from removal of both of its terminal hydroxy groups, which residue has a molecular weight of from 1,000 to 50,000 Daltons; w is an integer of from 1 to 3; and one of R₃ and R₄ is lower alkyl and the other is hydrogen or lower alkyl; and A is a hydrogen or an activated leaving group which when taken together with its attached oxygen atom forms an ester comprising, condensing a compound of the formula:



XX

wherein w, Y, R³, R⁴ and R⁵ are as above, Y is halide and R⁵ forms a hydrolyzable protecting group


with a compound of the formula:



V

wherein R, and PAG are as above, V is -OH or -NH₂,

to produce an ester of the formula:



XXI

wherein w, R, PAG, X, R³, R⁴ and R⁵ are as above

hydrolyzing said ester to form a free acid of the formula:




wherein R, PAG, X, R³, R⁴ and R⁵ are as above,

and reacting said free acid with a halide of an activated leaving group in the presence of a coupling agent to produce said activated ester.


54. (Original) The process of claim 53 wherein said leaving group is a N-hydroxysuccinimidyl group.

55. (Original) A process for producing an activated ester of the formula:




I-C

wherein R is hydrogen or lower alkyl, X is -O- or -NH-, A is a hydrogen or an activated leaving group which when taken together with its attached oxygen atom forms an ester, PAG¹ is a divalent residue of a polyalkylene glycol resulting from the removal of both of the terminal hydroxy groups, said residue having a molecular weight of from about 500 to about 25,000 Daltons, y is an integer from 0 to 3 and v is an integer from 1 to 3; and k is an integer from 1 to 2, comprising, condensing a compound of the formula:




wherein Y is halide, y and v are as above, and R⁵ forms a hydrolyzable ester protecting group

with a compound of the formula




wherein R, PAG¹ and k are as above, V is -OH or -NH₂,

to produce an ester of the formula:



wherein R, PAG¹, X, R⁵, k, v and y are as above,


hydrolyzing said ester to form a free acid of the formula:



wherein R, PAG¹, X, k, v and y are as above,
and reacting said free acid with a halide of an activated leaving group in the presence of a coupling agent to produce said activated ester.

56. (Original) The process of claim 55 wherein said leaving group is N-hydroxysuccinimidyl.

57. (Original) A conjugate of the formula




wherein P is a residue of a biopharmaceutical having a terminal hydroxy group wherein the terminal hydroxy group is removed, R, R₁ and R₂ are individually hydrogen or lower alkyl; X is -O- or -NH-; PAG is a divalent residue of polyalkylene glycol resulting from removal of both of its terminal hydroxy groups, which residue has a molecular weight of from 1,000 to 50,000 Daltons; n is an integer of from 0 to 1; and m is an integer of from 4 to 8.

58. (Original) The conjugate of claim 57 wherein P is a glycoside.

59. (Original) The conjugate of claim 58 wherein P is a residue of AZT.

60. (Original) The conjugate of claim 57 wherein X is -O-.
61. (Original) The conjugate of claim 60 wherein PAG is a polyethylene glycol residue having a molecular weight of 10,000 to 15,000.

62. (Original) A conjugate of the formula



wherein P is a residue of a biopharmaceutical having a terminal hydroxy group wherein the terminal hydroxy group is removed, R, R₁ and R₂ are individually hydrogen or lower alkyl; X is -O- or -NH-; PAG is a divalent residue of polyalkylene glycol resulting from removal of both of its terminal hydroxy groups, which residue has a molecular weight of from 1,000 to 50,000 Daltons; n is an integer of from 0 to 1; and m is an integer of from 4 to 8.

63. (Original) The conjugate of claim 62 wherein P is a residue of a protein or polypeptide.

64. (Original) The conjugate of claim 63 wherein X is -O-.

65. (Original) The conjugate of claim 64 wherein PAG is a polyethylene glycol residue having a molecular weight of about 10,000 to 15,000.

66. (Original) The conjugate of claim 63 wherein P is the polypeptide T-20 having a sequence according to SEQ ID NO: 1.

67. (Original) The conjugate of claim 64 wherein R is methyl.